Premier Health Anti-Asthmatic Monoclonal Antibody Faxed Order Form

Patient Name		_ Date of Birth
Patient MRN#	Patient's Allergies	
Ordering Physician	Physician's Phone/Fa	x #/
MVH Middletown: 513-974-5023 MVH Troy: 937-440-4503	Infusion Center Fax Numbe MVH North: 937-641-2378 MVH Greenville: 937-641-7205	<u>rs:</u> MVH South: 937-641-2676
-	SE HAVE PATIENT BRING CURREN	T MEDICATION LIST
Diagnosis: Severe Persiste	ent Asthma uncomplicated (145.50) 🗔 Mod	derate Persistent Asthma, uncomplicated (J45.40)
• —	ent Asthma w/ Acute Exacerbation (J45.51)	
_	thma (J82.83)	
PHYSICIAN ORDERS:		
PREMEDICATIONS:		
□		_
		-
	b please verify with the provider that the po ge anaphylactic reactions prior to schedulin	atient has a prescription for an epinephrine auto- Ig the patient.
omalizumab. If the patient did no provider for additional orders.	t bring their epinephrine autoinjector to the	when and how to use it prior to administration of ir appointment or if it is expired contact the
Xolair (Omalizumab) (Preferi	- /	
DOSE:	(Subcutaneous) FREQUENC	/:
	2 Hours after the first 3 injections, then 30r	ninutes for subsequent injections
Interleukin-5 Inhibitors		
Fasenra (Benralizumab) (Pref DOSE: 30mg Subcutaneous	ferred PH agent)	other :
🗌 Cinqair (Reslizumab)		
ADMINISTRATION: Doses will	Patient weight: admixed in a total volume of 50ml of 0.9% S ninutes, subsequent infusion can run over 20	
Nucala (Mepolizumab) DOSE: 100mg Subcutaneo	us q4weeks 🔲 other:	
Anti-Human Thymic Stromal Lyr	nphopoietin (Anti-TSLP) Agent	
Tezspire (Tezepelumab) DOSE: 210mg Subcutaneou	us q4weeks 🔲 other:	

Premier Health Anti-Asthmatic Monoclonal Antibody Faxed Order Form

Patient Name

Date of Birth____

ADDITIONAL ORDERS FOR BENRALIZUMAB, MEPOLIZUMAB, OMALIZUMAB & TEZEPELUMAB ADMINISTRATION

NURSING:

- \checkmark Vital signs baseline.
- $\sqrt{}$ Observe for hypersensitivity reactions after administration.
- $\sqrt{}$ Check vitals immediately after administration, then 30 minutes post administration.

Infusion Reaction Protocol:

 $\sqrt{}$ Premier Health standard infusion reaction protocols

ADDITIONAL ORDERS FOR RESILIZUMAB ADMINISTRATION

NURSING:

 $\sqrt{}$ Vital signs baseline, then 5 minutes after infusion has started then every 30 minutes.

 $\sqrt{}$ Observe for hypersensitivity reactions during infusion – this includes hypotension, shortness of breath, and rash.

 $\sqrt{}$ Check vitals immediately after completion, then 30 minutes post infusion. The patient must be observed for at least 30 minutes after completion of the iron infusion for hypersensitivity reactions.

 $\sqrt{}$ Discontinue IV and discharge patient upon completion of therapy.

Patient Name _____

Date of Birth_____

IV SALINE LOCK PANEL / CARRIER FLUID (for Reslizumab)

- √ Insert Saline Lock
- $\sqrt{}$ Saline Lock flush 10ml 0.9% NaCl as needed for line flush
- $\sqrt{}$ Discontinue Saline Lock on discharge

 $\sqrt{1000}$ NaCl 0.9% 1000 ml at 10 ml/hr Once PRN -- Admin Inst: : If infusion rate is less than 10 ml/hr or the infusion is a vesicant or a continuous IV solution is not infusing, a carrier fluid of 0.9% NaCl at 10 ml/hr may be initiated during infusion. DC Carrier fluid when infusion complete

Infusion Reaction Protocol:

 $\sqrt{}$ Premier Health standard infusion reaction protocols

Premier Health Anti-Asthmatic Monoclonal Antibody Faxed Order Form

Patient Name		Date of Birth			
Provider signature			-		
Printed provider name					
DateTi	me	_			
	_				
For Completion by Prior Authorization Team					
IF THE PATIENT HAS <u>INSURANCE OT</u> REQUIRED.	HER THAN MEDICARE OR TRA	ADITIONAL OHIO MEDICAID PRECEP	RTIFICATION IS		
PLEASE OBTAIN PRECERTIFICATION Precertification	AND INCLUDE AUTHORIZATIO	ON BELOW:			
	Date range:	# of infusion	s:		
□ No precertification necessary	-				
If no precert required, list name of whom you spoke with at insurance company and on what date					
Name:	Company:	Date:			